



Artery Research

ISSN (Online): 1876-4401

ISSN (Print): 1872-9312

Journal Home Page: <https://www.atlantis-press.com/journals/artres>

2.8: RELATIONSHIPS BETWEEN ADIPOSITY AND LEFT VENTRICULAR FUNCTION IN ADOLESCENTS: MEDIATION BY BLOOD PRESSURE AND OTHER CARDIOVASCULAR MEASURES

Hannah Taylor, Alun D. Hughes, Abigail Fraser, Laura Howe, George Davey Smith, Debbie Lawlor, Nishi Chaturvedi, Chloe Park

To cite this article: Hannah Taylor, Alun D. Hughes, Abigail Fraser, Laura Howe, George Davey Smith, Debbie Lawlor, Nishi Chaturvedi, Chloe Park (2018) 2.8: RELATIONSHIPS BETWEEN ADIPOSITY AND LEFT VENTRICULAR FUNCTION IN ADOLESCENTS: MEDIATION BY BLOOD PRESSURE AND OTHER CARDIOVASCULAR MEASURES, Artery Research 24:C, 71–72, DOI: <https://doi.org/10.1016/j.artres.2018.10.031>

To link to this article: <https://doi.org/10.1016/j.artres.2018.10.031>

Published online: 7 December 2019

keeping average wave speed in all terminal vessels constant ("proximal-stiffening", see Figure). An elastance heart model was applied at the inlet and simulations were performed with a one-dimensional flow solver (2).

Results: Proximal-stiffening and distal-stiffening had opposing effects on R but the same effects on mPP, whereas mPP increased monotonically with decreasing TAC and increasing Pfw in both settings (Figure).

Conclusion: Wave reflection per se does not provide protection from high mPP since greater reflection also entails greater transmitted pressure. Although a decreased R may accompany proximal arterial stiffening, the likely mechanism of increased mPP with aging is decreased TAC and greater Pfw.




Figure. Reflection coefficient, R (at every junction), TAC and Pfw vs mPP with proximal and distal stiffening. Arrows indicate effect of stiffening. c_{prox} and c_{dist} are wave speeds (m/s) in the proximal inlet segment and distal terminals (mean).

References

- Mitchell GF. Effects of central arterial aging on the structure and function of the peripheral vasculature: implications for end-organ damage. *J Appl Physiol.* 2008;105(5):1652-60.
- Mynard JP, Smolich JJ. One-Dimensional Haemodynamic Modeling and Wave Dynamics in the Entire Adult Circulation. *Ann Biomed Eng.* 2015;43(6):1443-60.

2.6

FEASIBILITY OF AORTIC WAVE INTENSITY ANALYSIS FROM SEQUENTIALLY ACQUIRED CARDIAC MRI AND NON-INVASIVE CENTRAL BLOOD PRESSURE

Anish Bhuvu^{1,2}, Niro Nadarajan³, Andrew D'Silva⁴, Camilla Torlasco⁵, Redha Boubertakh², Siana Jones³, Paul Scully^{3,2}, Rachel Bastiaenen⁴, Guy Lloyd², Sanjay Sharma⁴, James Moon^{3,2}, Kim Parker⁶, Charlotte Manisty^{3,2}, Alun Hughes^{3,7}

¹University College London, UK

²Barts Heart Centre, London, UK

³Institute of Cardiovascular Science, University College London, UK

⁴Cardiovascular Sciences Research Centre, St. George's, University of London, London, United Kingdom

⁵IRCCS, Istituto Auxologico Italiano, Milan, Italy

⁶Department of Bioengineering, Imperial College London, UK

⁷MRC Unit for Lifelong Health and Ageing at UCL, London, UK

Background: Wave intensity analysis (WIA) in the aorta offers important clinical and mechanistic insights but is difficult non-invasively. We performed WIA by combining high temporal resolution cardiovascular magnetic resonance (CMR) flow velocity and non-invasive central blood pressure (BP) waveform data.

Method: 206 healthy volunteers (36 ± 11 years, 47% male) underwent sequential phase contrast CMR (Siemens Aera 1.5T, 1.97×1.77 mm², ~ 9 ms temporal resolution) and supra-systolic oscillometric central BP (Uscom Ltd BP+) measurement. Velocity (U) and central pressure (P) waveforms (200 Hz) were aligned using the wave foot, and local wave speed was calculated both from the P-U slope during early systole (c) and the sum of squares method (cSS) (Figure 1), and compared with CMR aortic arch pulse wave velocity (PWV) by transit time.

Results: The peak intensity of the initial compression wave (di+1), backward compression wave (di-) and protodiastolic decompression wave (di+2) were 69.5 ± 28 , -6.6 ± 4.2 and 6.2 ± 2.5 W/m² respectively. PWV correlated with c or cSS ($r = 0.60$, and 0.68 respectively; bias -1.3 [limits of agreement: -3.8 to 1.2 m/s], and bias -0.64 [limits of agreement: -3.0 to 1.7 m/s] respectively), Figure 1.

Conclusion: Wave intensity patterns and values are similar to those measured using invasive methods. Local wave speed showed good agreement with PWV. CMR and central blood pressure provides a novel non-invasive technique for performing wave intensity analysis and is feasible for large scale studies.




Figure 1. Calculation of wave speed from a pressure – velocity (P-U) loop and comparison with pulse wave velocity by transit time. Top left: alignment of scaled pressure (blue) and velocity (red) waveforms and example of wave intensity analysis showing initial compression (di+1), backward compression (di-) and protodiastolic decompression (di+2) waves. Top right: P-U loop showing wave speed measurement in early systole (c); and using sum of squares (cSS). Bottom left and Bottom right: Correlation and Bland-Altman analysis of cSS and PWV from phase-contrast MRI showing good correlation and slight underestimation.

2.7

FITNESS MODIFIES THE ASSOCIATION BETWEEN EXERCISE BLOOD PRESSURE AND LEFT-VENTRICULAR MASS IN ADOLESCENCE

Zhengzheng Huang¹, Ricardo Fonseca², James Sharman², Nish Chaturvedi³, George Smith⁴, Deborah Lawlor⁴, Laura Howe⁴, Chloe Park³, Alun Hughes³, Martin Schultz², Martin Schultz²

¹Menzies Institute for Medical Research, Hobart, Australia

²Menzies Institute for Medical Research, University of Tasmania, Hobart, Australia

³Institute of Cardiovascular Science, University College London, London, UK

⁴MRC Integrative Epidemiology Unit, University of Bristol, Bristol, UK

Objective: Exaggerated exercise blood pressure (BP) is associated with higher left-ventricular mass index (LVMI). Paradoxically, exercise BP and LVMI may be higher with greater fitness, but underlying factors are poorly understood. This study aimed to determine the influence of fitness on exercise BP and its relationship with LVMI in adolescents.

Methods: 4835 adolescents from the Avon Longitudinal Study of Parents and Children, aged 15.4(0.3) years, 49% male completed a submaximal cycle test. Exercise BP was measured immediately on test cessation and fitness calculated as physical work capacity 170 adjusted for lean body-mass. LVMI ($n = 1589$), cardiac output (CO, $n = 1628$) and total peripheral resistance (TPR, $n = 1628$) were measured by echocardiography 2.4 (0.4) years later.

Results: Each unit of fitness was associated with a 6.46 mmHg increase (95% CI: 5.83, 7.09) in exercise systolic BP. Exercise systolic BP increased step-wise by third of fitness (difference 6.06 mmHg, 95% CI: 4.99, 7.13 first vs. middle; 11.13 mmHg, 10.05, 12.20 middle vs. highest). Each 5 mmHg increase in exercise systolic BP was associated with 0.25 g/m². 7 (0.16–0.35) greater LVMI, attenuated with adjustment for fitness. There was evidence of an interaction between fitness and exercise BP on LVMI, more-marked in the middle fitness third (difference -0.27 g/m². 7, -0.51 , 0.04 vs. first third), but similar in lowest and highest fitness thirds. CO increased (difference 0.06 L/min, -0.05 , 0.17 ; 0.23 L/min, 0.12, 0.34), TPR decreased (difference -0.13 AU, -0.84 , 0.59 ; -1.08 AU, -0.180 , -0.35 with fitness).

Conclusion: Fitness may modify associations between exercise BP and LVMI in adolescence. Higher CO, but lower TPR suggests a physiological exercise BP-LVMI relationship with higher fitness, rather than pathological elevations in exercise BP and LVMI.

2.8

RELATIONSHIPS BETWEEN ADIPOSITY AND LEFT VENTRICULAR FUNCTION IN ADOLESCENTS: MEDIATION BY BLOOD PRESSURE AND OTHER CARDIOVASCULAR MEASURES

Hannah Taylor¹, Alun D. Hughes^{2,3}, Abigail Fraser⁴, Laura Howe⁴, George Davey Smith⁴, Debbie Lawlor⁴, Nishi Chaturvedi^{3,2}, Chloe Park²

¹University College London, London, UK

²Department of Population Science & Experimental Medicine, Institute of Cardiovascular Science, University College London, UK

³MRC Unit for Lifelong Health and Ageing at UCL, University College London, UK

⁴MRC Integrative Epidemiology Unit, University of Bristol, UK

Introduction: Increased adiposity is associated with poorer left ventricular (LV) function but the mediating role of blood pressure (BP) and other cardiovascular measures is unknown. We investigated the roles of potential mediators in adolescents in a UK birth cohort, the Avon Longitudinal Study of Parents and Children.

Methods: Fat and lean mass were assessed by dual energy X-ray absorptiometry (DXA) in 2,068 individuals (age 17.7(SD 0.32) years; 45% male; weight 67(SD 13) kg). BP was measured and echocardiography performed. Tissue Doppler systolic (s'), early diastolic longitudinal function (e') and transmural E/A were used as measures of LV systolic and diastolic function respectively. Mediation was estimated using structural equation modelling; height, sex, socioeconomic position, lean mass and smoking were included as potential confounders in all models.

Results: Table 1 shows the total and direct (total – indirect (mediated)) effects (standardized β) for each potential mediator, alongside %mediation. No convincing associations were found between fat mass and s' or e' .

Conclusions: MAP, HR and EDV, an indicator of preload, are important mediators of the effect of adiposity on diastolic function in adolescence, while TPR appears to account for none of the effect. Absence of mediation by PP, in combination with the mediation by TAC, suggests that stroke volume contributes some mediative effect. These findings emphasise the importance of adiposity and risk factor control in adolescence.

Table 1. Mediation of the association between fat mass and E/A

Mediator	Direct effect (standardized β)	Total effect (standardized β)	% mediation
None		-0.15 (95%CI -0.20, -0.10)	
Mean arterial pressure (MAP)	-0.11	-0.14	21
Pulse pressure (PP)	-0.13	-0.13	0
Heart rate (HR)	-0.11	-0.14	21
Total arterial elastance (TAE)	-0.15	-0.14	-7
Total peripheral resistance (TPR)	-0.15	-0.15	0
End-diastolic volume (EDV)	-0.17	-0.15	-13

Oral Session III – Clinical Aspects, Hypertension & Diabetes

3.1

STRAIN DISCONTINUITIES IN CAROTID ATHEROSCLEROTIC PLAQUES – A NOVEL MARKER FOR PLAQUE VULNERABILITY?

Tim Vonk^{1,2}, Evelien Hermeling¹, Floris Schreuder³, Werner Mess¹, Eline Kooi¹

¹Maastricht University Medical Center, Maastricht, the Netherlands

²Zuyderland Medical Center, Heerlen, the Netherlands

³Radboud University Medical Center, Nijmegen, the Netherlands

Objective: Rupture of atherosclerotic plaques is ultimately a biomechanical event. We aim to develop and validate a novel method using ultrasound radiofrequency (RF) measurements to determine intraplaque inhomogeneities in the strain distribution with a high axial resolution which may identify rupture-prone plaques.

Methods: Ultrasound examinations were performed on CCAs and ICAs of 17 patients with recent stroke/TIA and 5 asymptomatic patients (39 CCAs and ICAs). Strain distribution was computed from RF data with a depth resolution <0.5 mm. In the plaque-free CCA, radial wall strain was compared with distension for validation. Two observers analyzed the data. In the ICAs, strain inhomogeneities in the plaque were studied.

Results: Within the CCA wall, the strain (observer 1: $-7.4 \pm 2.7\%$, observer 2: $-6.4 \pm 2.6\%$) had good intra-subject precision (1.6%), accuracy (correlation with relative distension, observer 1: $r = 0.69$; $p < 0.0001$, observer 2: $r = 0.68$; $p < 0.0001$) and intra-observer variability (ICC of 0.681). In the ICA, strain inhomogeneities demarcated in 58% of cases the plaque-arteria boundary. The percentage of ICAs with ≥ 1 strain inhomogeneities was increasing with the degree of stenosis ($p = 0.03$, 95%-CI). Strain inhomogeneities were more frequent in ICAs at the symptomatic side of stroke/TIA (odds-ratio = 4.7; $p = 0.07$). Strain in the deviating strain regions was higher at the symptomatic compared to the asymptomatic side ($p = 0.02$).

Conclusion: The proposed method to assess local radial strain distribution proved to be accurate and precise. In ICA, areas with high strain inhomogeneities were more frequent in symptomatic plaques vs asymptomatic

plaques. These strain inhomogeneities could be a promising novel marker for plaque vulnerability.




Figure 1. Examples of strain distribution in carotid plaques

Crude B-mode (echo, left panels) and corresponding strain distribution in the ICA (right panels); blood flow direction is from right to left. Note that in some cases the strain distribution is rather homogeneous (A), whereas in other cases a more heterogeneous distribution of strain is observed (B-D). Narrow but elongated regions with highly deviating strain (indicated by the white arrows) are often found near the boundary between the plaque and the supporting adventitia layer (B and D) and can appear either as diastolic-to-systolic wall compression (B) or as wall tension (C and D).

3.2

RESERVOIR PRESSURE INTEGRAL IS INDEPENDENTLY ASSOCIATED WITH THE REDUCTION IN RENAL FUNCTION IN AN OLDER POPULATION

Kunihiko Aizawa¹, Francesco Casanova¹, David M. Mawson², Kim M. Gooding¹, W. David Strain¹, Phillip E. Gates¹, Gerd Östling³, Faisal Khan⁴, Helen M. Colhoun⁵, Carlo Palombo⁶, Kim H. Parker⁷, Jan Nilsson³, Angela C. Shore¹, Alun D. Hughes⁸

¹Diabetes and Vascular Medicine Research Centre, NIHR Exeter Clinical Research Facility, University of Exeter Medical School, Exeter, UK

²Diabetes and Vascular Medicine Research Centre, NIHR Exeter Clinical Research Facility, University of Exeter Medical School, Exeter, UK

³Department of Clinical Sciences, Lund University, Malmö, Sweden

⁴Division of Molecular & Clinical Medicine, University of Dundee, Dundee, UK

⁵Centre for Genomic and Experimental Medicine, University of Edinburgh, Edinburgh, UK

⁶Department of Surgical, Medical, Molecular and Critical Area Pathology, University of Pisa, Pisa, Italy

⁷Department of Bioengineering, Imperial College, London, UK

⁸Institute of Cardiovascular Science, University College London, London, UK

Background: Central haemodynamic parameters derived from reservoir pressure analysis (RPA-parameters) exhibit prognostic utility. Alterations in reservoir function could have an unfavourable influence on target organs, such as the kidneys. We determined in older adults whether these RPA-parameters would be associated with the reduction in estimated glomerular filtration rate (eGFR) during a 3-year follow-up period.

Methods: We studied 544 individuals (69.4 ± 7.9 yrs, 195F, 235CVD) at baseline and after 3 years. RPA-parameters including reservoir pressure integral (INTPR), peak reservoir pressure, excess pressure integral, systolic and diastolic rate constants were obtained by radial artery tonometry.

Results: After 3 years, 95 individuals (72.4 ± 7.6 yrs, 26F) had an eGFR reduction of more than $5 \text{ ml/min/1.73 m}^2/\text{year}$. A multivariate logistic regression analysis revealed that baseline INTPR was independently associated with the eGFR reduction after adjusting for conventional risk factors and baseline renal function [odds ratio 0.975 (95% CI, 0.958–0.993), $p < 0.01$]. When the eGFR reduction was expressed as a continuous variable, baseline INTPR was also independently associated with changes in eGFR ($\beta = 0.115$, $p < 0.01$; multivariate linear regression with adjustment for